

GERASIMOV-DRELL-HEARN AND BURKHARDT-COTTINGHAM SUM RULES FOR PROTON, NEUTRON AND DEUTERON

O.V. Teryaev

*Bogoliubov Laboratory of Theoretical Physics
Joint Institute for Nuclear Research, Dubna
141980, Russia*¹

Abstract

A description of the generalized Gerasimov-Drell-Hearn sum rules for proton, neutron and deuteron is suggested. The proton consideration confirms the earlier conjecture, that the structure function g_T features a smooth Q^2 -dependence, while the structure function g_2 is changing rapidly, due to the elastic contribution to the Burkhardt-Cottingham sum rule. The neutron case features the close connection to the Bjorken sum rule. The deuteron case reveals the crucial role of photodesintegration channel, resulting in the strong nuclear correction at very low Q^2 . The new approximate sum rules for this channel reflecting the delicate interplay between nucleon and nuclear structure are suggested.

Key-words: Sum Rules, QCD, Photodesintegration, Binding

1 Introduction

The generalized (Q^2 -dependent) Gerasimov-Drell-Hearn (GDH) sum rules provide the important link between the low energy description of hadrons by classical GDH [1, 2] sum rules for real photons ($Q^2 = 0$) and PQCD description at large Q^2 . They are just being tested experimentally with a high accuracy for both proton and neutron [3, 4, 5]. The striking feature of the proton data is the low ($\sim 200 - 250 MeV^2$) “crossover” point, which is in complete agreement with the prediction [6, 7], published almost 10 years ago. It was based on the relation to another fundamental sum rule, namely, Burkhardt-Cottingham (BC) sum rule for structure function g_2 . Its elastic contribution is the main source of strong Q^2 -dependence, while the contribution of the other structure function, $g_T = g_1 + g_2$ is smooth. In the present report I review this approach for proton, as well as more recent treatment of neutron using the connection to Bjorken sum rule. Also, following the general line of this conference, exploring the nuclear aspects of QCD, I will address the problem of (generalized) GDH and BC sum rule for deuteron. I will use the approximation of zero binding energy and show, that the crucial role in the description of nuclear effects is played by the photodesintegration channel. Its effect is manifested at rather low Q of the order of pion mass and leads to the set of the new approximate “finite-energy” sum rules for photodesintegration channel.

¹E-mail: teryaev@thsun1.jinr.ru

2 Proton: Burkhardt-Cottingham sum rule for longitudinally polarized particles [6, 7]

Let us consider the Q^2 -dependent integral

$$I_1(Q^2) = \frac{2M^2}{Q^2} \int_0^1 g_1(x, Q^2) dx. \quad (1)$$

It is defined for *all* Q^2 , leaving no ambiguity for the choice of "virtual photon flux factor", and g_1 is the obvious generalization for all Q^2 of the standard scale-invariant $g_1(x)$. Therefore, the comparison with its another definitions should be made with some care, although the ambiguity due to the choice of smooth function I_T (see below) makes this discrepancy actually beyond the accuracy of our approach. Note that the elastic contribution at $x = 1$ is not included in the above sum rule. One recovers then at $Q^2 = 0$ the GDH sum rule

$$I_1(0) = -\frac{\mu_A^2}{4} \quad (2)$$

where μ_A is the nucleon anomalous magnetic moment in nuclear magnetons. While $I_1(0)$ is always negative, its value at large Q^2 is determined by the Q^2 independent integral $\int_0^1 g_1(x) dx$, which is positive for the proton and negative for the neutron.

The separation of the contributions of g_T and g_2 leads to the decomposition of I_1 as the difference of the contributions of two form factors I_T and I_2

$$I_1 = I_T - I_2, \quad (3)$$

where

$$I_T(Q^2) = \frac{2M^2}{Q^2} \int_0^1 g_T(x, Q^2) dx, \quad I_2(Q^2) = \frac{2M^2}{Q^2} \int_0^1 g_2(x, Q^2) dx; \quad (4)$$

$$g_T = g_1 + g_2. \quad (5)$$

There are solid theoretical arguments to expect a strong Q^2 -dependence of I_2 . It is the well-known Burkhardt-Cottingham sum rule [8], derived independently by Schwinger [9], using a rather different method. It states that

$$I_2(Q^2) = \frac{1}{4} \mu G_M(Q^2) \frac{\mu G_M(Q^2) - G_E(Q^2)}{1 + \frac{Q^2}{4M^2}}, \quad (6)$$

where μ is the nucleon magnetic moment, G 's denoting the familiar Sachs form factors which are dimensionless and normalized to unity at $Q^2 = 0$. For large Q^2 one can neglect the r.h.s. of (6) and get

$$\int_0^1 g_2(x) dx = 0. \quad (7)$$

In particular, from Eq.(7) it follows that

$$I_2(0) = \frac{\mu_A^2 + \mu_A e}{4}, \quad (8)$$

e being the nucleon charge in elementary units. To reproduce the GDH value (see Eq.(3)) one should have

$$I_T(0) = \frac{\mu_A e}{4}, \quad (9)$$

which was indeed proved by Schwinger [9]. The importance of the g_2 contribution can already be seen, as the entire μ_A - term for the GDH sum rule is provided by I_2 .

Note that I_T does not differ from I_1 for large Q^2 due to the BC sum rule, but it is *positive* in the proton case. It is possible to obtain a smooth interpolation for $I_T^p(Q^2)$ between large Q^2 and $Q^2 = 0$ [6].

$$I_T^p(Q^2) = \theta(Q_0^2 - Q^2) \left(\frac{\mu_A}{4} - \frac{2M^2 Q^2}{(Q_0^2)^2} \Gamma_1^p \right) + \theta(Q^2 - Q_0^2) \frac{2M^2}{Q^2} \Gamma_1^p, \quad (10)$$

where $\Gamma_1^p = \int_0^1 g_1^p(x) dx$. The continuity of the function and of its derivative is guaranteed with the choice $Q_0^2 = (16M^2/\mu_A) \Gamma_1^p \sim 1 \text{ GeV}^2$, where the integral is given by the world average proton data. It is quite reasonable to distinguish the perturbative and the non-perturbative regions. As a result one obtains a crossing point at $Q^2 \sim 0.2 \text{ GeV}^2$, below the resonance region [6], while the asymptotic behaviour for $Q^2 > 1 \text{ GeV}^2$ and positive value at $Q^2 = 0.5 \text{ GeV}^2$ is in a good agreement with the HERMES [5], E143 [3] and JLAB [4] data. This smooth interpolation seems to be very reasonable in the framework of the QCD sum rules method as well. Then one should choose some "dominant" tensor structure to study the Q^2 -dependence of its scalar coefficient and T_1 appears to be a good candidate. This seems also promising from another point of view. It is not trivial to obtain within the QCD sum rules approach the GDH value at $Q^2 = 0$. Since the r.h.s. of (12) is linear in μ_A , it may be possible to obtain it using the Ward identities, just like the normalization condition for the pion form factor. This, in turn opens the possibility to apply the powerful tool of quark-hadron duality. The latter relies on the perturbative theory, and it is quite clear that it is much more plausible to describe linear (one-loop), than quadratic terms. One should recall here that while sum rule for I_T was checked in QED long ago [10], the GDH sum rule required much more efforts [11], as it gets non-trivial contributions only at two loops order, although zero appearing at one loop level is also non-trivial [12, 13].

Note also that large contribution of g_2 by no mean contradicts the resonance approaches [14] and may be considered complimentary to them. The central role here plays the $\Delta(1232)$: it provides a significant amount of GDH integral at $Q^2 = 0$ and gives a clear qualitative explanation of rapid Q^2 -dependence [15]. The Δ photoproduction is dominated by the magnetic dipole form factor, leading to a negative I_1 . The sign change is just related to the fast decrease of the Δ contribution. In order to compare this picture with our approach we have separated [7] the Δ contribution to I_T and I_2 . To do this we just calculated the photoproduction Born diagram using the well-known expressions for the covariant form factors G_M , G_E and G_C . The resulting expression is rather lengthy but it has a remarkable property: the leading G_M^2 term contains *only* the tensor T_2 at any Q^2 . This fact is confirmed if one performs the contraction with the virtual photon density matrix: in particular, if one takes the standard definition with the kinematical zero at $Q^2 = 0$, the result should be attributed to g_1 . In approach [7] the nonzero g_1 is due to the absence of T_1 since $g_1 = -g_2$.

3 Neutron: link to Bjorken sum rule [16]

To generalize this approach to the neutron case, one needs a similar smooth parameterization of g_T for the neutron. Since the value at $Q^2 = 0$ is equal to zero, it is not sufficient to limit oneself to the simplest linear parameterization. One needs to add a term, quadratic in Q^2 . A simple parameterization providing the continuity of the function and its derivative was suggested in [17], which however, leads to the result being in contradiction with the

data. This does not seem to be occasional, bearing in mind the argument presented above. Indeed, the general reason supporting the smoothness of interpolation for g_T is its linearity in μ_A . As soon as this term appears to be equal to zero for some occasional reason (which, in our case, is nothing else than the neutrality of neutron!), there is also no more reason to expect the smoothness as well.

To bypass this obstacle, we use the difference between the proton and neutron instead of neutron itself. Although it is possible, in principle, to construct the smooth interpolation for functions g_1 themselves [15], it also does not fit the suggested general argument, as r.h.s. is proportional to $\mu_{A,p}^2 - \mu_{A,n}^2$, so that it is quadratic and, moreover, has an additional suppression due to the smallness of isoscalar anomalous magnetic moment. Indeed, the smooth approximation for g_1 resulted in the $Q^2 \sim 5GeV^2$, leading to the rather early scaling violations, do not observed experimentally. Instead, it was recently suggested the following parameterization for isovector formfactor I_T^{p-n} :

$$I_T^{p-n}(Q^2) = \theta(Q_1^2 - Q^2) \left(\frac{\mu_A}{4} - \frac{2M^2 Q^2}{(Q_1^2)^2} \Gamma_1^{p-n} \right) + \theta(Q^2 - Q_1^2) \frac{2M^2}{Q^2} \Gamma_1^{p-n}, \quad (11)$$

where transition value Q_1^2 may be determined by the continuity conditions in a similar way, and we get value $Q_1^2 \sim 1.3GeV^2$, being of the same order as in the proton case.

The elastic contribution to BC sum rule should be included for neutron separately, so we need the neutron elastic form factors. While the electric one, might be neglected, the magnetic form factor is well described by the dipole formula [?]

$$G_M(Q^2) = \frac{1}{(1 + Q^2/0.71)^2}. \quad (12)$$

As a result [16], the neutron g_1 stays quite close to its asymptotic value until rather low Q are achieved. Moreover, the experimental data happen to bear some traces of the structure, generated in our approach by the interplay of Q_0 and Q_1 .

4 Deuteron: sum of squares vs square of sum and new finite energy sum rules for photodesintegration channel

The deuteron case poses the new problem, which is manifested already for real photon scattering ($Q^2 = 0$). Considering deuteron as a fundamental particle of spin 1 one get the deuteron anomalous magnetic moment, which is very small due to the approximate cancellation of proton and neutron AMM, in the r.h.s. of (2).

At the same time, for large energies (and/or Q^2) one should expect, due to the small binding energy to get the incoherent sum of the proton and neutron contributions which is rather large.

To make this effect even more dramatic one may consider the following *gedankenexperiment*. Let the binding to be adiabatically switched on. Than, for zero binding, one should get the *sum of squares* of the proton and neutron AMM, while for *arbitrary* weak binding it should become equal to the *square of their sum*, as weakness of binding will make the deviation from AMM additivity negligible.

So, what transforms the sum of squares to square of sums? The answer is almost obvious. For arbitrary small binding there is a new channel, photodesintegration $d \rightarrow p + n$,

absent for zero binding, when it corresponds to elastic contribution do not entering to GDH integral.

As soon as binding energy is small, one may assume that the inelastic contributions (starting from the pion threshold) for free proton and neutron are not affected by binding. If so, the inelastic contribution to deuteron GDH integral is equal to the sum of the ones for free proton and neutron. As a result, one may get the following finite energy sum rule:

$$\int_{\nu_{PD}}^{\nu_{\pi}} \frac{d\nu}{\nu} \Delta\sigma_d(\nu) = -\frac{4\pi^2\alpha}{M^2} ((\sum \mu_A)^2 - \sum \mu_A^2) \approx -\frac{8\pi^2}{M^2} \mu_A^P \mu_A^n \quad (13)$$

Here ν_{PD} and ν_{π} are the photodesintegration and inelastic (one-pion) thresholds, respectively. The accuracy of this sum rule is guaranteed by smallness of deuteron binding energy ϵ . The relevant small parameter may be estimated as its ratio β to the minimal energy, which should be transferred to nucleon to produce the pion,

$$\beta = \frac{2M\epsilon}{m_{\pi}^2} \sim 10\%. \quad (14)$$

The contribution of photodesintegration channel was first estimated long ago by S.B. Gerasimov [18], but the use of non-relativistic approximation resulted in the correction to (13) of about 50%, so that the significant role of nuclear effects at high-energies was expected. However, the more elaborate recent relativistic calculation [19] led to the validity of (13) with the accuracy of about 10%, which is the one expected from the small binding energy approximation (SBEA) suggested above.

This approximation should become much more accurate in the virtual photon case, as now

$$\beta = \frac{2M\epsilon}{m_{\pi}^2 + Q^2}.$$

This mean that it should provide very good description down to $Q^2 \sim m_{\pi}^2$ (recall, that even for $Q^2 \sim 0$ the accuracy of SBEA is still rather good, so that it is m_{π}^2 which determines the characteristic value of Q^2).

Consequently, down to low $Q^2 \sim m_{\pi}^2$ the GDH integral should be very close to the sum of the proton and neutron values. The difference between "square of sum" and "sum of squares" (which, in turn is rapidly decreasing because of the decrease of the form factors) should be provided by the photodesintegration channel contribution, corresponding to very large $x_B \sim 1 - 2Mm_{\pi}/Q^2$ and therefore also small.

So, it is convenient to decompose the generalized GDH SR and consider the photodesintegration and inelastic channels separately. While the inelastic channel contribution should be close to the sum of proton and neutron down to real photon limit, while the photodesintegration one should rapidly grow at low Q^2 and tend to the value (13).

One can immediately make more quantitative predictions for BC SR, valid for all Q^2 . Like in the case of generalized GDH SR, SBEA should be good at all Q^2 , resulting in another sum rule for photodesintegration channel.

$$\int_{x_{\pi}}^{x_{PD}} g_2^d(x) dx = \frac{1}{4} ((\sum \mu G_M)^2 - \sum \mu G_M \sum G_E - \sum \mu G_M (\mu G_M - G_E)), \quad (15)$$

where x_{π}, x_{PD} correspond to inelastic and photodesintegration channels, respectively. This sum rule may be probably checked at JLAB, while (13) may be studied at TUNL [20].

I am indebted to J. Soffer for the fruitful collaboration in this field and to S.B. Gerasimov and A. Schäfer for valuable comments. This investigation was supported by INTAS(Contract 00-00587)

References

- [1] S. B. Gerasimov, Yad. Fiz. **2**, 598(1965) [Sov. J. Nucl Phys. **2**, 430(1966)].
- [2] S. D. Drell and A. C. Hearn, Phys. Rev. Lett. **16**, 908 (1966).
- [3] E143Collaboration, K. Abe et al., Phys. Rev. Lett.**78**, 815 (1997).
- [4] M. Amarian *et al.*, arXiv:nucl-ex/0205020.
- [5] A. Airapetian *et al.* [HERMES Collaboration], Eur. Phys. J. C **26**, 527 (2003)
- [6] J. Soffer and O. Teryaev, Phys. Rev. Lett. **70**, 3373 (1993).
- [7] J. Soffer and O. Teryaev, Phys. Rev. **D51**, 25 (1995).
- [8] H. Burkhardt and W. N. Cottingham, Ann. Phys. (N.Y.) **16**, 543 (1970).
- [9] J. Schwinger, Proc. Nat. Acad. Sci. U.S.A. **72**, 1559 (1975).
- [10] Wu-Yang Tsai, L.DeRaad and K.A.Milton, Phys. Rev. **D11**, 3537(1975).
- [11] D.A. Dicus and R. Vega, Phys. Lett. **B501**, 44 (2001)
- [12] S.B. Gerasimov and J. Moulin, JINR E2-6722, 1972. G. Altarelli, N. Cabibbo and L. Maiani, Phys. Lett. **B40**, 415 (1972)
- [13] S.J. Brodsky and I. Schmidt, Phys. Lett. **B351**, 344 (1995)
- [14] V. D. Burkert and B. L. Ioffe, Phys. Lett. **B296**, 223 (1992), J. Exp. Theor. Phys. **78**, 619 (1994) [Zh. Eksp. Teor. Fiz. **105**, 1153 (1994)]; V. D. Burkert, arXiv:nucl-ex/0109004.
- [15] B. L. Ioffe, V. A. Khoze, L. N. Lipatov, *Hard Processes*, (North-Holland,Amsterdam, 1984).
- [16] J. Soffer and O. V. Teryaev, Phys. Lett. B **545**, 323 (2002)
- [17] J. Soffer and O. Teryaev, Phys. Rev. **D56**, 7458 (1997).
- [18] S.B. Gerasimov, Yad. Fiz. **5**, 1263 (1967)
- [19] H. Arenhövel, nucl-th/0006083
- [20] H.R. Weller, Published in *Mainz 2000, Gerasimov-Drell-Hearn sum rule and the nucleon spin structure in the resonance region* 145-156